



SCIENTIFIC PANEL
ON RESPONSIBLE PLANT NUTRITION

Issue Brief 06 – February 2024

Fertilizer and soil health for enhanced productivity and sustainability in sub-Saharan Africa

by the Scientific Panel on Responsible Plant Nutrition



SCIENTIFIC PANEL
ON RESPONSIBLE PLANT NUTRITION

Contents

- 3 Key points
- 5 What is soil health?
- 8 What are the issues?
- 19 Solutions and needed actions
- 24 What would success look like?
- 26 References
- 28 Authors, citation and contact



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

Key points

1. Soil health indicators and targets: Soil health is commonly defined in relation to achieving sufficient crop yields while maintaining the future productive capacity of soils and the ecosystem services soils provide. Soil organic matter is a key soil health indicator, along with available P and K, soil acidity, and water-stable aggregation. There is no consensus yet on easily measurable, meaningful biological indicators.

2. Towards fertilizer and soil health: Managing soil health in sub-Saharan Africa is complex:

- To bolster agricultural productivity, both fertilizers and organic inputs are essential. Together with sound agronomic practices, superior seed varieties, soil amendments, and other measures, this forms the basis of Integrated Soil Fertility Management.
- On healthier soils, the efficiencies of both fertilizers and water tend to be higher. Hence, enhancing soil quality directly contributes to improved agricultural performance.
- Increased input of plant nutrients as fertilizer is indispensable for sustainable agriculture that seeks to increase food production and restore soil health. Relying solely on organic inputs often results in soil nutrient depletion and degradation.
- Building soil health and rehabilitating degraded soils are long-term processes; in stark contrast to the rapid depletion of soil organic matter and nutrients caused by practices such as land clearing or excessive nutrient removal.

3. Addressing climate change requires choices: Climate change affects the relationship between fertilizer and soil health. Support that addresses specific risks created by climate change will be essential for farmers to engage in sustainable fertilizer and soil health management practices. Such investments are likely to have a high return.

4. Enabling farmers: Bottlenecks in engaging smallholder farmers in soil health-restoring practices are large upfront costs, the length of time required to achieve visible benefits, and access to affordable inputs. Fertilizer subsidies need to be re-designed with respect to soil health outcomes. Soil health is a public good, thus justifying greater public investment in it.

5. Soil health investments require localized actions: Soil health management should always be localized and context-based. The specific solutions for building soil health depend on soil characteristics, environment, the farming system, crops grown, and socioeconomic abilities.



SCIENTIFIC PANEL
ON RESPONSIBLE PLANT NUTRITION

SOIL HEALTH FRAMEWORK




Figure 1. Adapted model of the soil health framework from (3). Arrows present processes connecting the system components and management strategies. The blue line relates to water availability and the green lines indicate direct impacts of mineral fertilizers on biomass/yield and SOC.



SCIENTIFIC PANEL
ON RESPONSIBLE PLANT NUTRITION

What is soil health?

Broadly speaking, soil health refers to the overall condition and quality of soil, including its physical, chemical, and biological characteristics that define the capacity of a soil to function as a vital living ecosystem that sustains plants, animals, and humans (2).



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

Healthy soil has good water-holding capacity, drainage, and resistance to erosion, a balanced pH as well as adequate levels of plant nutrients and organic matter. Moreover, healthy soil sustains a diverse community of soil organisms, such as bacteria, fungi, and fauna, which play important roles in nutrient cycling and maintaining or building soil structure. Based on these properties, soil health directly influences the productivity, resilience, and sustainability of agricultural and natural ecosystems.

Measuring soil health involves assessing a range of physical, chemical, and biological properties that contribute to soil fertility, productivity, and sustainability. Specific indicators describe soil health, depending on the context and management goals of the particular system:

- **Physical indicators:** These include soil texture, structure, porosity, water-holding capacity, and infiltration rate. Physical indicators can be measured using soil cores or pits, or instruments such as penetrometers, tensiometers, and soil moisture sensors.
- **Chemical indicators:** These include soil pH, nutrient levels (e.g., nitrogen, phosphorus, potassium, micronutrients), organic matter content (Box 1), and cation exchange capacity. To measure chemical indicators, soil samples are analyzed in a laboratory, or using the crop nutrient status as an indicator of soil nutrient supplying capacity.
- **Biological indicators:** These include the presence and diversity of soil organisms, such as bacteria, fungi, earthworms, or nematodes. Biological indicators can be measured using techniques such as soil enzyme assays, microbial biomass measurements, and DNA analysis. Their interpretation in relation to specific soil functions of agronomic importance is, however, still challenging.

Box 1: Soil organic matter

In this document, we refer to soil organic carbon (SOC), noting that soil organic matter (SOM) is constituted of organic compounds that contain C, N, and other elements. In that sense, SOC is a measurable indicator of SOM. Since the current document refers both to SOM as a property that hosts several functions and to SOC as an element that can be managed by soil management practices, we decided to refer only to SOC in the text, noting that the amount of SOC and SOM is strongly correlated in most circumstances.

Combined, these indicators give a good idea on the health status of the soil. Several tools and protocols are available for quantifying soil health, including the Comprehensive Assessment of Soil Health (20), the soil management assessment framework (21), regional guidelines for Europe (22), or simplified national programs such as the Soil Health Card program in India. Generally speaking, validation of these tools in SSA has been limited.

The use of specific indicators and thresholds to quantify soil health depends on the functions that a soil is envisaged to support (23). From an agronomic viewpoint, healthy soils increase crop yields by releasing nutrients, supporting water retention, facilitating root growth, and protecting crops from pests and diseases (24). The content of organic carbon in the soil is of cross-cutting importance to its health status, as soil organic matter favorably affects all these properties.



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

The timely and sufficient availability of plant nutrients is particularly critical as most of them impact crop performance as well as soil health. For example, depending on the soil type, complex processes immobilize P in the soil and may thus hamper its availability. In SSA, exchangeable K is too low in many soils to achieve good yields of root and tuber crops. Soil acidity is another key indicator; crop growth is commonly limited at pH values below 5 and above 8. Physical indicators of soil health are generally related to the ease of root growth through the top- and subsoil and the regulation of water infiltration and storage. While it would be important to quantify or assess their impact on soil health, current microbial measurements are not easy to interpret and may not necessarily provide credible inferences about soil health status (25). That said, progress has been made in Brazil, where soil bioanalysis (SoilBio) has been increasingly used by farmers (Box 2).

In addition to the need of specific indicators to assess soil health, three other aspects must be considered: (i) spatial heterogeneity of some soil properties and the need to integrate this into protocols aiming to measure change in specific indicators; (ii) the rate of change of some indicators, which can be relatively fast (e.g., changes in available P as affected by P fertilizer application) or slow, especially when these are measured against a relatively large background (e.g., changes in SOC); and (iii) the cost of measuring indicators over large areas of land, acknowledging that remote and proximal soil spectral methods have considerably reduced the cost while retaining accuracy for, e.g., assessment of SOC and total N (30). Alternatively, instead of directly measuring soil health indicators, one could assess the change in crop and soil management practices as related to expected changes in the status of key indicators through empirical or mechanistic modeling.

Crop productivity directly depends on key indicators of soil health such as SOC, available P, exchangeable K, soil pH, or water-stable aggregation. In this Brief, we focus more on soil organic carbon (SOC) because it is a central indicator of soil health and challenging to manage, whereas solutions to address low available P, K, S, micro-nutrients, and acid soils exist.

Box 2: The Brazilian experience with indicators of soil health

The economy in Brazil is strongly based on agriculture, so rebuilding soil fertility and maintaining soil quality/health are priority goals. For decades, chemical and physical parameters, including SOC, were evaluated searching for indicators of soil quality/health that are reliable and feasible for large-scale analysis. In the late 1990s, microbial parameters of biomass C and N and a metabolic quotient were suggested as interesting indicators; more effective and able to detect changes earlier than any chemical or physical parameters (26). In a long-term study in the Cerrado region, a savannah region with many similarities to SSA, microbial indicators were interpreted as a function of the relative cumulative yields of maize and soybean and SOC, and the critical levels were defined based on criteria similar to those used in soil nutrient calibration tests (27). Subsequent studies found that only two enzymes, arylsulfatase and β -glucosidase, were associated with the carbon and sulfur cycles, respectively (28). These two soil enzymes, in addition to routine chemical analyses, were integrated into a Soil Bioanalysis technology in 2020 (29), which has been exponentially used by farmers to assess soil health. Consistent results obtained in two decades of studies in Brazil show that it may take years until changes in SOC are detected, whereas faster changes can be detected in microbial enzymatic activities.



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

What are the issues?

1. Fertilizer use and crop productivity increase, but progress is diverse and slow when aggregated for sub-Saharan Africa

Agricultural production in sub-Saharan Africa (SSA) needs to urgently increase to ensure food and nutrition security of the growing population (1). If intensification is not successful and massive cropland expansion was to be avoided, SSA would become even more dependent on imports of cereals and other staples than it is already. In 2004, the then-UN-general secretary Kofi Annan called for a uniquely African Green Revolution to take root within the rich diversity of the continent. In 2006, the African Union, through the *Abuja Declaration on Fertilizer for an African Green Revolution*, recognized the critical need to enhance access to fertilizers to achieve such an African Green Revolution, given the poverty trap caused by poor and declining soil fertility. Yet, progress has been slow (2, 3) and the problem remains, often resulting in a per capita yield decline for many countries in SSA. Although average fertilizer nutrients use in SSA was only 15 kg N+P₂O₅+K₂O/ha in 2022 (excluding South Africa), several countries have approached or actually reached the Abuja target of 50 kg/ha. Concerning is, however, that average fertilizer consumption in SSA has stagnated since 2019.



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

In Asia, agricultural production rose fast and was primarily driven by yield growth. Average cereal yields have more than tripled in the past 60 years, whereas cereal area only increased by about 20%. In Africa, on the other hand, crop yield growth has been slow whereas the harvested crop area more than doubled during the same period (Figure 2, left). Notwithstanding disappointing continent-wide figures, in recent years few countries in Africa have started to move toward a trajectory similar to that of Asia. In Ethiopia, for example, maize yields nearly doubled in the past 20 years, mostly on existing land (Figure 2, right). Increased fertilizer, better varieties and other improvements drove that change.




Figure 2. Relative changes in cereal grain yield and harvested area in Asia and Africa (Left) and maize yield and area in Sub-Saharan Africa (SSA), Ethiopia, Nigeria and Rwanda (Right), 1961-2020. Data shown are 5-year averages, with the average of 1961-1965 set as 100. Source: [FAOSTAT](#).

Maintaining soil health is a globally relevant issue, but the situation in SSA is unique because large areas under smallholder farming continue to suffer from nutrient mining, eroding the productive capacity of large areas of agricultural soils. Sub-Saharan Africa remains the only region with negative nutrient balances that continue to increase over time, for instance for nutrients such as P or K (Figure 3). Actual nutrient balances are even more negative because losses due to removed crop residues, leaching, runoff, or erosion are not included in Figure 3. It must be noted, however, that significant differences exist among regions and countries in Africa, with West and Central Africa generally faring worst.

It is now widely accepted that producing crops in that manner -- at the cost of soil health -- is an unsustainable poverty trap. Increasing crop productivity to meet current and future food needs and improving soil health in SSA will only be successful with the increased and efficient use of fertilizer nutrients (4). A second African Fertilizer and Soil Health Summit is scheduled to take place in 2024 to draw up new plans for action.



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION




Figure 3. Cropland phosphorus (left) and potassium (right) input-output balances for different sub-regions of Africa, 1961-2021. Source: FAO Cropland Nutrient Balance database (5).

2. From focussing on increasing crop yields to recognizing soil health as a prerequisite for sustainability

In the 1980s and 1990s, in SSA increasing attention was directed to the needs of small-scale farms. Work on nutrient mining revealed widespread soil nutrient deficits, highlighting the critical role of soil fertility (6, 7). To address smallholder farms and low-fertility soil, a soil management paradigm was articulated (8) that highlighted the need to 'rely more on biological processes to optimize nutrient cycling, minimize external inputs and maximize the efficiency of their use.' The initial focus was on using crop varieties that were adapted to adverse soil conditions, such as soil acidity and poor nutrient availability, and increasing the availability of organic inputs to complement fertilizer, with a strong emphasis on improved (leguminous) fallows or intercrops. Soil organic matter and the combined use of mineral fertilizers and organic inputs were emphasized as means to increase efficiencies of nutrient use.

Concurrently, so-called low-external-input sustainable agriculture (LEISA) received much attention (9), although it

also became clear that the low yields achieved without nutrient inputs could not address the critical issues of poverty and food security. Due to these concerns, the focus shifted toward replenishing soil fertility (10).

The Integrated Soil Fertility Management (ISFM) paradigm (11) emerged in the 2000s, based on decades of soil and crop research in Africa. In contrast to earlier paradigms, ISFM recognizes the critical need to use all organic and mineral nutrient resources efficiently and focus on using fertilizers as entry points to intensifying smallholder agriculture. It follows a stepwise approach to rehabilitate, improve, and rebuild soils. First, fertilizers and improved crop varieties are applied to raise productivity and rebuild the fertility base of the soil. The next step is to incorporate organic inputs into soil management to rebuild the soil organic matter pool, which is key to soil health and integral to multiple soil functions. For example, legume inoculation with N-fixing bacteria (rhizobia) can be used to improve legume yield and thus the in-situ availability of organic inputs in the form of crop residues (12). There is scope to deploy various bio-stimulants to increase crop yield and residues (13), though care needs to be taken since many ineffective bio-stimulants are currently available in SSA (14). Other amendments or practices such as lime application, water harvesting, or deep tillage, are also included to address local constraints.



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

Over the past two decades, major investments were also made to promote conservation agriculture, largely driven by concerns about soil erosion and loss. Smallholder farmers face, however, a number of challenges to implement the commonly recommended three pillars of conservation agriculture (15): (i) minimizing soil disturbance through tillage; (ii) maintaining a continuous soil cover; and (iii) diversifying crops. To ensure sufficient crop production, a fourth principle for conservation agriculture was proposed: using mineral fertilizer appropriately (16).

More recently, ‘agroecology’ and ‘regenerative agriculture’ have received increasing attention, which consider other ecosystem services besides crop productivity as equally important. Soil health is viewed as a key indicator for the delivery of those other services. Regenerative agriculture favors practices that (i) encourage water infiltration and prevent soil erosion; (ii) build soil organic carbon (SOC) and enhance biological nutrient cycling; (iii) foster plant diversity; (iv) integrate livestock; and (v) reduce reliance on external inputs (17). Applying these ideas to the typical farming situations found in SSA faces, however, major shortcomings (18) because plant nutrients first need to be made available in order to reach many other goals.

In summary, the Green Revolution has allowed increased food production to exceed global population growth over the past decades, except in SSA. As a region, SSA experienced many decades in which agricultural concepts and practices were promoted that neither benefitted soil health nor farmers and food consumers. It is now urgent to correct that and focus on the things that must come first to sustainably intensify agriculture, while also recognizing the increasing awareness of environmental and societal challenges associated with food systems (4). Managing nutrients for soil, crop, and human health is of highest priority for SSA.



SCIENTIFIC PANEL


ON RESPONSIBLE PLANT NUTRITION

and other structures enhance the volume of soil that can be penetrated by roots.

The nutrient pool and its dynamics vary for different nutrients and can be affected by agronomic practices and edaphic processes. For example, crop residues and manure can be left on the land surface or incorporated, serving as substrates for (micro)organisms and thus raising the carbon pool of the soil (32). Plants obtain their nutrients from the available pool of nutrients, influenced by the cation exchange capacity (CEC) of soils. Clay minerals primarily determine CEC, with clay soils having a greater capacity to hold nutrients than sandy soils. In soils with low CEC, SOC is a key supplier of CEC. Also, soil acidity, soil water content for nutrient dissolution and diffusion, and mineral and organic fertilizer application affect nutrient pool size and availability. Nitrogen fertilizers can induce soil acidity, based on their content of ammonium.

Using fertilizers is essential to increase crop yields and the amounts of organic residues, thus raising SOC and sequestering N, P, and S. As seen in Figure 5, on average, a 1.3% annual increase in crop grain yields was associated with a 0.4% annual increase in SOC stock across Africa, Asia, and Latin America.


Figure 5. Relative annual changes in crop productivity and soil organic carbon stock (over 0-20 cm) (%) after changes in land management practices (33). Crop species (symbols): B, beans; C, cassava; M, maize; P, sweet potatoes; R, rice; S, soybean; s, sorghum; W, wheat. Regions: Africa (black); Asia (green); and Latin America (blue). The solid line is the standard major axis regression for all data points.





SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION



The large scatter indicates that many other factors besides increased crop yields affect changes in SOC. Hence, additional practices are needed to ensure that increased crop yields and biomass production improve soil health.

Whereas long-term application of fertilizers in temperate climates increases or maintains SOC (34), research on the impact of fertilizer use on SOC in different cropping systems in SSA is scant. General indications suggest that common soil health-promoting practices are not increasing SOC (35). By reviewing the effect of fertilizer on SOC, Beillouin et al. (36) found only one meta-analysis covering experiments in SSA (37), which showed no significant effects of fertilizer on SOC (Figure 6). Yet, only a few of the included studies were carried out for five or more years. There are indications, however, that including soybean can improve SOC content and land productivity (38).

Figure 6. Synthesis of meta-analyses evaluating the effect of fertilizers on SOC. Points represent the mean effect size, and error bars represent confidence intervals. Numbers represent paired data used in a given meta-analysis. N: nitrogen; P: phosphorus; and K: potassium. Data from the evidence map of Beillouin et al. (36).



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION




Figure 7. Boxplot of annual change in SOC between the end and the start of the long-term experiment in the treatment with fertilizer (+F) and the control (-F) for 25 long-term experiments in sub-Saharan Africa. Source: M. Corbeels et al., unpublished.

In line with the meta-analysis findings of Chivenge et al. (37), a review of 25 long-term experiments conducted in SSA with contrasting climate, soils, and mineral fertilizer inputs revealed no clear evidence that SOC decreased less strongly over time upon treatment with a mineral fertilizer (Figure 7). Probably, lower crop yield levels (and thus C inputs) in SSA explain the different trends observed in croplands globally. That said, even in long-term experiments in temperate regions with fairly high rates of fertilizer use, the impact on SOC is modest – but almost always positive (39). Another way to increase SOC is through biomass transfer, i.e. applying organic inputs that have been produced outside the plot. This may, however, not represent a net benefit if it leads to depletion of soil health on the land on which this biomass was produced.

4. Limitations to increasing soil organic carbon

To what extent can SOC be increased in reality and under the conditions of SSA? Soil organic carbon under natural vegetation could be viewed as a proxy for what could be achieved in a given soil and climate combination. However, while the native SOC may provide some general guidance for the C sequestration potential of soils, it is not a realistic goal for SOC restoration under agriculture. Natural ecosystems are not meant to produce and export much biomass, but are dominated by plant species that allocate much of their carbon to roots, which are primary sources for forming stable soil organic matter. As such, attainable SOC restoration under agriculture may reach only 60-70% of its natural potential (40).




Figure 8. Changes in soil organic C following clearing of woodland, as affected by duration and intensity of farming at Chikwaka, Zimbabwe(41).

Decomposing organic material releases large amounts of CO₂, such as after land clearing. Hence, through continuous cultivation and with limited to no mineral or organic inputs, the carbon content of the soil continues to decline. That said, even soils under commercial agricultural practices do not restore SOC to the levels under natural vegetation (Figure 8). Whether this observation is a major issue or not depends on the amount of SOC that is required to maintain critical ecosystem functions for the intended land use, as affected by soil properties and climatic conditions -- a question that is still awaiting a conclusive answer. One could also argue that the function of soil and its impact on the ecosystem has changed as land management changed from natural vegetation to agriculture and therefore a lower SOC became acceptable.

Increasing SOC contents is an option for soils in Africa that have the capacity to stabilize soil organic matter, such as clay soils with high oxide contents that are common in large areas of Africa. African soils can be important C sinks since they have experienced a massive decline in SOC, mainly as a result of low C inputs into soil. Applying organic inputs can increase SOC, yet limited availability of such organic inputs is a constraint. Adding organic matter from external sources is limited in scope and may lead to C depletion in the areas where this organic matter was collected. Hence, the application of fertilizer is a more realistic path towards increasing SOC through growing larger amounts of biomass in situ, particularly roots. It should, however, also be noted that per ton of carbon about 100 kg of N, 10 kg of P and 10 kg of S are required to form stable SOM (42).



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

5. Economic realities

Although fertilizers are key components of sustainable food systems, their use in Africa heavily depends on precarious politics. Fertilizer use varies greatly between and within countries due to different micro- and macroeconomic conditions. Moreover, biophysical conditions, such as the amount of rainfall and soil type, are important; the risk of crop failure resulting from low rainfall is a strong disincentive to the purchase and use of fertilizers on subsistence crops. This section focuses on the economic dimensions of fertilizer use.

Across SSA, crop yields are much lower than what is attainable under the given environmental conditions and available technologies. Yield gaps in smallholder agriculture in SSA are the largest in the world (43) and low use of fertilizers combined with low fertilizer use efficiency are a major cause of that (44). Closing the yield gap in SSA is the primary route for improving food security and the well-being of rural households (1), while farmers in SSA will need to use much more fertilizer than they currently use.

However, even though it may be technically feasible to increase crop yields in many regions of SSA, it is not always economically viable for farmers because of relatively high fertilizer prices and relatively low crop produce prices. Location-specific ecological and economic conditions greatly affect crop response to fertilizer and economic returns on fertilizer investments (45). The latter study showed that (i) the average economic yield gap (the difference between current yield and profit-maximizing yield) was about 25% of the





SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION



ecological yield gap; (ii) though maize yields could be profitably doubled, the economic incentives to do so are weak due to unfavorable input:output price ratios; and (iii) risk from variable seasonal rainfall makes this worse. Considerable variation of fertilizer profitability exists within countries. These findings emphasize the need for tailoring fertilizer recommendations to smaller areas based on the local environmental and economic conditions.

Variations in agricultural technology adoption can be attributed to differences between farmers, farms, and the environments in which they operate. Farmers are less likely to apply fertilizers if the risk of non-response or non-profitable response is high, fertilizer prices are too high (and there is no government support), or crop prices are too low.

Land is another key factor for crop production. Farmers with large farm sizes and land ownership are likely to invest in fertilizers and soil health practices. They are also likely to have significant or adequate amounts of organic matter resources. Farmers without secure land tenure are less likely to invest in fertilizers or other soil health practices than those who have more certainty to reap the medium to long-term benefits (46–48). Farmers who own land are also able to access credit as a pre-requisite for investing in technologies. A farmer's age, gender (49), and education level are other mediating factors affecting the uptake of technologies and correspond to the farmer's knowledge, ability, preferences, and risk appetite. While this is an important topic, directly affecting the use of fertilizer and its impact on soil health, a detailed analysis of the factors underlying this topic is out of scope of the current paper.



SCIENTIFIC PANEL
ON RESPONSIBLE PLANT NUTRITION



Solutions and needed actions

A number of solutions and actions are required to ensure that fertilizers are used efficiently and their application results in increased crop yields, while soil health is maintained or even improved.



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

1. Consensus on what constitutes soil health, based on specific functions and environments

A key performance indicator framework is required that defines soil health and the key functions of a healthy soil for the intended land use. Such indicators need to be measurable and interpretable in relation to the key soil functions, as well as to the management practices affecting them. Soil organic carbon is generally acknowledged as a key indicator. Besides SOC, available P, exchangeable K, soil acidity, and water-stable aggregation could complete the group of key indicators for soil health in relation to crop productivity in SSA. Currently, there is no consensus on easily measurable biological indicators, notwithstanding progress being made elsewhere (Box 2). Where necessary, additional soil or crop indicators may be used to quantify soil supplying capacity of specific plant nutrients, including micronutrients.

2. Monitoring schemes to assess progress and support soil health management

National or regional monitoring schemes should support public investments in soil health by monitoring change over time and space. Such soil information should be openly available, thus also improving digital soil information resources across Africa (50). Another important route is soil health monitoring as part of vertically integrated supply chains, for example in relation to sustainable production standards implemented by companies. Harmonized and scientifically sound soil health assessment frameworks need to integrate globally accepted standard operational procedures and data management systems, implemented by public extension agents as well as private sector agronomists.

3. Raising awareness on how farmers and society benefit from improved soil health

Soil health benefits society as a whole by providing and regulating numerous ecosystem services. Smallholder farmers benefit from the health of soil as crop yields increase and production systems become more efficient, ultimately leading to better returns on investment. Soil health is also a public good. Hence, investing in soil health, informed by actual quantification of these benefits, means ultimately investing in public health. Soil health only changes meaningfully over longer periods of time. Hence, longer-term incentive schemes are needed, partly supported by public funding, to enable farmers to invest in soil health management. In this same context, it is also important to make the public aware that well-managed fertilizers do not ‘kill’ soil — a misconception regularly reaching public discussion fora in SSA — but are actually essential for good soil health.

4. Technical, data-driven solutions to improve soil health beyond beliefs and wishful thinking

Improving soil health is a laudable and necessary goal of an agricultural transformation agenda, yet not always easily achievable. For example, soil health improvement is limited by poor nutrient retention in sandy soils. Similarly, specific agronomic practices are assumed to improve soil health, e.g., as is often the case for conservation agriculture, but they are not always universally applicable. Evidence often suggests that gaining soil health depends on many other factors that need to be embedded in improved, locally adapted practices.



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

5. Include investments in fertilizer and soil health in climate adaptation or mitigation

Climate change affects the relationship between fertilizer and soil health. De-risking agriculture will be essential for farmers to engage in fertilizer and soil health management practices while addressing specific hazards created by climate change. Investments in soil health improvement also contribute to increase farming system resilience through adaptation and mitigation.

6. Implement policies that enable farmers to invest in soil health

Land tenure is key, but farmers in SSA will also continue to require fertilizer subsidies since there are no alternative sources to provide the amount and diversity of nutrients needed to grow crops. Redesigning fertilizer subsidies is crucial with respect to soil health management. Such subsidies need to be designed smartly, and include educating farmers on how to maximize the use efficiency of the input and how to manage locally available organic resources and crop residues in combination with fertilizers.





SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

Who needs to do what?

The global fertilizer industry needs to integrate aspects of soil health management in fertilizer recommendations and ensure that the right fertilizer formulations are available for the right crops and soils to maximize crop biomass production. Co-supply of fertilizers and effective bio-stimulants could provide an entry point for the increased use of the latter.

Service providers need to integrate knowledge on soil health management practices into agronomic advisory services and share the information efficiently with farmers.

Scientists need to identify and characterize key indicators for soil health, and practical protocols and tools for rapid and efficient monitoring of soil health status. They should further characterize agronomic practices that restore or maintain soil health under different environment and management conditions. Soil health monitoring systems need to be set up to capture changes in soil health conditions as related practices move to scale.

Policy makers need to facilitate farmers to invest in soil health improvement, which is a longer-term objective that requires public support, including redesigned fertilizer subsidies. Lessons learnt from countries that released policies related to payment for SOC sequestration in soil could form a basis for developing similar policies in SSA. Organic waste stream management and use can be another route towards increasing the availability of organic inputs, especially in rural areas near urban centers. National and/or regional soil health monitoring units with the necessary expertise and tools need to be established and maintained.



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

Who needs to do what?

Investors need to integrate soil health management as a key condition for financial support and support the implementation of soil health-promoting financial mechanisms. Sustaining public support to facilitate farmers' investments in soil health will be required in the medium to long-term to build soil health to reasonable levels.

Civil society organizations need to raise awareness across all sectors of society on the importance of soil health, the role of nutrient inputs, both organic and mineral, as well as its maintenance and improvement. They need to show how improving soil health can counteract climate change-related stresses, yield decline, and environmental degradation. Their work should be based on scientific evidence, not ideology or beliefs.

The retail sector needs to include soil health-positive production practices in their processes for certifying responsible production.

Consumers are encouraged to buy soil health-positive food and actively engage in discussions related to soil health, as is currently done in relation to climate change.



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

What would success look like?

Access to soil health management information and practice is facilitated

Integrated Soil Fertility Management (ISFM) is mainstreamed into agricultural development programs. Fertilizer subsidy schemes include support for soil health management practices in the context of ISFM, including aspects of 4R for fertilizer management. Funders provide medium to long-term support to soil health improvement in their respective agricultural development programs. At least 70% of smallholder farmers on the continent are accessing targeted agronomic recommendations for specific crops, soils, and climatic conditions to ensure greater efficiency and sustainable use of fertilizers.

A more diverse range of fertilizer products and organic inputs is available and affordable

The domestic production and distribution of fertilizer and organic inputs is tripled within the next 10 years thus enabling access and affordability for smallholder farmers. To improve production, procurement, and distribution of fertilizer and organic inputs, the Africa Fertilizer Financing Mechanism (AFFM) is fully operational. Policies and regulations are formulated and implemented to create a conducive environment for fertilizer and soil health interventions.

Soil health conditions are improving as an outcome of the African Fertilizer and Soil Health Summit of 2024

Consensus is reached on key indicators related to soil health, standard operational procedures for measuring these together with trends or thresholds indicating positive change. Upon uptake of soil health management practices, many smallholder farming systems in SSA register positive change at scale. The agronomic efficiency of fertilizer is doubled (e.g., from 10 to 20 kg maize grain per kg fertilizer nutrient) with marked increases in crop yields and crop residue availability, thus also increasing the in-situ availability of organic inputs for improving soil health. At least 30% of degraded soil is restored within 10 years.



What would success look like?

Advisory and extension services integrate soil health management information

Advisory services are up to date about fertilizer and soil health management, supported by a large network of certified agronomists providing bundled services to farming communities. Feedback on the performance of agronomic practices informs the R&D community to further improve the efficacy and efficiency of specific recommendations.

National and regional science capacity is capable of generating and disseminating locally relevant soil health management recommendations

Science capacity is strengthened at the national and regional level, especially in areas such as data science, advanced analytics, and artificial intelligence, thus supporting the development, validation, and dissemination of site-specific, local recommendations. Effective linkages are established between the R&D community, service providers, and last-mile delivery staff.

Public awareness and support

The public is aware of the role of soil health in addressing many of the challenges faced by farmers and by society as a whole, and appreciates the role of both fertilizer and organic nutrient inputs in maintaining and improving soil health. The retail sector is adopting soil health-positive certification schemes and consumers are preferably buying soil health-positive products.



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

References

1. M. K. van Ittersum et al., Can sub-Saharan Africa feed itself? *Proc. Natl. Acad. Sci.* 113, 14964–14969 (2016).
2. N. Winnie et al., Assessment of the 2006 Abuja Fertilizer Declaration with emphasis on nitrogen use efficiency to reduce yield gaps in maize production. *Front. Sustain. Food Syst.* 5 (2022).
3. B. Vanlauwe et al., Fertilizer and soil health in Africa: The role of fertilizer in building soil health to sustain farming and address climate change (2023) .
4. B. Vanlauwe, A. Dobermann, Sustainable intensification of agriculture in sub-Saharan Africa: first things first! *Front. Agr. Sci. Eng.* 7, 376–382 (2020).
5. C. I. Ludemann et al., A global reference database in FAOSTAT of cropland nutrient budgets and nutrient use efficiency: nitrogen, phosphorus and potassium, 1961–2020. *Earth Syst. Sci. Data* (2023).
6. J. J. Stoorvogel, E. M. A. Smaling, B. H. Janssen, Calculating soil nutrient balances in Africa at different scales. 1. Supra-national scale. *Fert. Res.* 35, 227–235 (1993).
7. E. M. A. Smaling, J. J. Stoorvogel, P. N. Windmeijer, Calculating soil nutrient balances in Africa at different scales. 2. District scale. *Fert. Res.* 35, 237–250 (1993).
8. P. A. Sánchez, in *Transactions of the 15th World Congress of Soil Science, Vol. 1, ISSS, Ed. (International Soc. Soil Sci. and Mexican Soc. Soil Sci., Acapulco, 1994), vol. 1, pp. 65–88.*
9. C. Reijntjes, B. Haverkort, A. Waters-Bayer, *Farming for the future: An introduction to Low-External Input and Sustainable Agriculture* (Macmillan, London, 1992).
10. R. J. Buresh, P. A. Sanchez, F. Calhoun, *Replenishing soil fertility in Africa. SSSA Special Publication Number 51* (Soil Science Society of America, American Society of Agronomy, Madison, Wisconsin, USA, 1997).
11. B. Vanlauwe et al., Integrated soil fertility management: Operational definition and consequences for implementation and dissemination. *Outlook on Agriculture.* 39, 17–24 (2010).
12. K. E. Giller, F. Kanampiu, M. Hungria, B. Vanlauwe, The role of nitrogen fixation in African smallholder agriculture. *Agriculture, Ecosystems & Environment.* 285, 106601 (2019).
13. C. N. Sible, J. R. Seebauer, F. E. Below, Plant biostimulants: a categorical review, their implications for row crop production, and relation to soil health indicators. *Agronomy.* 11, 1297 (2021).
14. C. Masso, J. R. Awuor Ochieng, B. Vanlauwe, Worldwide contrast in application of bio-fertilizers for sustainable agriculture: lessons for Sub-Saharan Africa. *Journal of Biology, Agriculture and Healthcare.* 5, 34–50 (2015).
15. K. E. Giller, E. Witter, M. Corbeels, P. Tittonell, Conservation agriculture and smallholder farming in Africa: The heretics' view. *Field Crops Res.* 114, 23–34 (2009).
16. B. Vanlauwe et al., A fourth principle is required to define Conservation Agriculture in sub-Saharan Africa: The appropriate use of fertilizer to enhance crop productivity. *Field Crops Res.* 155, 10–13 (2014).
17. K. E. Giller, R. Hijbeek, J. A. Andersson, J. Sumberg, Regenerative Agriculture: An agronomic perspective. *Outlook Agric.* 003072702199806 (2021).
18. K. E. Giller, Why the buzz on regenerative agriculture? *Growing Africa.* 1, 12–16 (2022).
19. T. Chang, G. Feng, V. Paul, A. Adeli, J. P. Brooks, Soil health assessment methods: Progress, applications and comparison. *Adv. Agronomy.* 172, 129–210 (2022).
20. B. N. Moebius-Clune et al., Comprehensive assessment of soil health. The Cornell framework (Cornell University, Ithaca, New York, ed. 3, 2022).



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

References

21. S. S. Andrews, D. L. Karlen, C. A. Cambardella, The soil management assessment framework. A quantitative soil quality evaluation method. *Soil Science Society of America Journal*. 68, 1945–1962 (2004).
22. EEA, Soil monitoring in Europe. Indicators and thresholds for soil quality assessments (Publications Office of the European Union, Luxembourg, 2023).
23. H.-J. Vogel et al., Quantitative evaluation of soil functions: potential and state. *Front. Environ. Sci.* 7 (2019).
24. S. Kuyah et al., Innovative agronomic practices for sustainable intensification in sub-Saharan Africa. A review. *Agron. Sustain. Dev.* 41, 1–21 (2021).
25. N. Fierer, S. A. Wood, C. P. Bueno de Mesquita, How microbes can, and cannot, be used to assess soil health. *Soil Biol. Biochem.* 153, 108111 (2021).
26. E. L. Balota, A. Colozzi-Filho, D. S. Andrade, M. Hungria, Biomassa microbiana e sua atividade em solos sob diferentes sistemas de preparo e sucessão de culturas. *Rev. Bras. Ciênc. Solo.* 22, 641–649 (1998).
27. Alves de Castro Lopes, André et al., Interpretation of microbial soil indicators as a function of crop yield and organic carbon. *Soil Sci. Soc. Am. J.* 77, 461–472 (2013).
28. I. d. C. Mendes et al., Critical limits for microbial indicators in tropical Oxisols at post-harvest: The FERTBIO soil sample concept. *Applied Soil Ecology*. 139, 85–93 (2019).
29. I. C. Mendes et al., in *Soil health and sustainable agriculture in Brazil*, Madison: Soil Science Society of America, I. C. Mendes, M. R. Cherubin, Eds. (Soil Science Society of America, Madison, WI, 2024), pp. 220–250.
30. K. D. Shepherd et al., A global soil spectral calibration library and estimation service. *Soil Security*. 7, 100061 (2022).
31. A. Bationo et al., in *Soil and fertilizers: managing the environmental footprint*, R. Lal, Ed. (CRC Press, Boca Raton, FL, 2020), pp. 67–102.
32. A. Bationo, A. Buerkert, in *Managing organic matter in tropical soils: scope and limitations*, C. Martius, H. Tiessen, P. L. G. Vlek, Eds. (Springer, Dordrecht, 2001), pp. 131–142.
33. J.-F. Soussana et al., Matching policy and science: Rationale for the ‘4 per 1000 - soils for food security and climate’ initiative. *Soil and Tillage Research*. 188, 3–15 (2019).
34. R. J. Haynes, R. Naidu, Influence of lime, fertilizer and manure applications on soil organic matter content and soil physical conditions: a review. *Nutr Cycl Agroecosyst.* 51, 123–137 (1998).
35. J. Kihara, P. Bolo, M. Kinyua, S. S. Nyawira, R. Sommer, *Soil health and ecosystem services: Lessons from sub-Saharan Africa (SSA)*. *Geoderma*. 370, 114342 (2020).
36. D. Beillouin et al., A global overview of studies about land management, land-use change, and climate change effects on soil organic carbon. *Global Change Biol.* 28, 1690–1702 (2022).
37. P. Chivenge, B. Vanlauwe, J. Six, Does the combined application of organic and mineral nutrient sources influence maize productivity? A meta-analysis. *Plant Soil*. 342, 1–30 (2011).
38. J. B. Naab, G. Y. Mahama, I. Yahaya, P. V. V. Prasad, Conservation agriculture improves soil quality, crop yield, and incomes of smallholder farmers in North Western Ghana. *Front. Plant Sci.* 8, 996 (2017).
39. J. K. Ladha, C. K. Reddy, A. T. Padre, C. van Kessel, Role of nitrogen fertilization in sustaining organic matter in cultivated soils. *J. Environ. Qual.* 40, 1756–1766 (2011).
40. D. S. Powlson, P. R. Poulton, M. J. Glendining, A. J. Macdonald, K. W. Goulding, Is it possible to attain the same soil organic matter content in arable agricultural soils as under natural vegetation? *Outlook Agric.* 51, 91–104 (2022).



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

References

41. S. Zingore, C. Manyame, P. Nyamugafata, K. E. Giller, Long-term changes in organic matter of woodland soils cleared for arable cropping in Zimbabwe. *European Journal of Soil Science*. 56, 727–736 (2005).
42. C. A. Kirkby, A. E. Richardson, L. J. Wade, M. Conyers, J. A. Kirkegaard, Inorganic nutrients increase humification efficiency and C-sequestration in an annually cropped soil. *PLoS ONE*. 11, e0153698 (2016).
43. P. Tittonell, K. E. Giller, When yield gaps are poverty traps: The paradigm of ecological intensification in African smallholder agriculture. *Field Crops Res*. 143, 76–90 (2013).
44. H. F. M. ten Berge et al., Maize crop nutrient input requirements for food security in sub-Saharan Africa. *Global Food Security*. 23, 9–21 (2019).
45. C. Bonilla-Cedrez, J. Chamberlin, R. J. Hijmans, Fertilizer and grain prices constrain food production in sub-Saharan Africa. *Nature Food*. 2, 766–772 (2021).
46. A. Abdulai, V. Owusu, R. Goetz, Land tenure differences and investment in land improvement measures: Theoretical and empirical analyses. *J. Dev. Econ*. 96, 66–78 (2011).
47. S. Bedeke, W. Vanhove, M. Gezahegn, K. Natarajan, P. van Damme, Adoption of climate change adaptation strategies by maize-dependent smallholders in Ethiopia. *NJAS: Wageningen Journal of Life Sciences*. 88, 96–104 (2019).
48. R. H. Oostendorp, F. Zaal, Land acquisition and the adoption of soil and water conservation techniques: a duration analysis for Kenya and The Philippines. *World Development*. 40, 1240–1254 (2012).
49. A. Hirpa Tufa et al., Gender differences in technology adoption and agricultural productivity: Evidence from Malawi. *World Development*. 159, 106027 (2022).
50. T. Hengl et al., African soil properties and nutrients mapped at 30 m spatial resolution using two-scale ensemble machine learning. *Sci. Rep*. 11, 6130 (2021).

Authors, citation and contact

Authors: Members of the Scientific Panel on Responsible Plant Nutrition.

Bernard Vanlauwe (International Institute of Tropical Agriculture, Kenya), Patrick H Brown (University of California - Davis, USA), Tom Bruulsema (Plant Nutrition Canada), Achim Dobermann (International Fertilizer Association, France), Mariangela Hungria (Embrapa, Brazil), Kaushik Majumdar (African Plant Nutrition Institute, Morocco), Michael McLaughlin (University of Adelaide, Australia), Claudia Wagner-Riddle (University of Guelph, Canada), Xin Zhang (University of Maryland, USA), Fang-jie Zhao (Nanjing Agricultural University, China).

Citation: Scientific Panel on Responsible Plant Nutrition. 2024. Fertilizer and soil health for enhanced productivity and sustainability in sub-Saharan Africa. Issue Brief 06. Available at <https://sprpn.org>.

Acknowledgement: This brief is based on a technical background document, prepared for the African Fertilizer and Soil Health Summit, held in Nairobi in 2024.

Further information: Scientific Panel on Responsible Plant Nutrition, info@sprpn.org