



SCIENTIFIC PANEL  
ON RESPONSIBLE PLANT NUTRITION

Issue Brief 07 – December 2024

# Mitigating greenhouse gas emissions through responsible plant nutrition

by the Scientific Panel on Responsible Plant Nutrition



# Contents

- 3 Key points
- 5 What is the Issue?
- 8 What is happening?
- 11 Interventions to reduce N<sub>2</sub>O emissions
- 16 Interventions to increase yields and N use efficiency
- 18 What can be done?
- 24 Who needs to do what?
- 26 What would success look like?
- 28 References
- 31 Authors, citation and contact



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

# Key points

The provision of nutrients for crop growth in the form of inorganic or organic fertilizers results in greenhouse gas (GHG) emissions from the processes involved in their manufacture and delivery to the farm as well as from the soil after application.

Opportunities exist for reducing emissions along the whole fertilizer supply chain. At issue is how to achieve emissions reductions at large scales in balance with the need to secure food production and farmers' incomes. Reducing soil GHG emissions in the form of nitrous oxide (N<sub>2</sub>O) and improving fertilizer nitrogen use efficiency (NUE) are of particular importance.


Emissions of N<sub>2</sub>O from the soil are highly variable, driven by soil and climatic factors as well as the interaction of those factors with management practices. Despite this variability a range of interventions under the framework of responsible plant nutrition have been shown to reduce emissions. These interventions include both 4R nutrient stewardship practices and crop and soil management practices that enhance crop yields and crop nitrogen removal. The 4R source practices that inhibit nitrogen transformations can directly reduce

field emissions of N<sub>2</sub>O, while all 4R components (source, rate, time, and place) along with crop and soil management can improve NUE, with impact on all emissions from the whole fertilizer supply chain. These practices have potential to mitigate a large fraction of annual emissions of GHG from fertilizer use within the next few decades, particularly when applied in combination.

We suggest actions that would prioritize the most appropriate mechanisms in target regions to ensure short-term success while also allowing for long-term progress. Social and economic constraints affecting farmer capacity to adopt emission-mitigating practices need to be addressed to achieve broader impact. New emission inventory methodologies and market-driven credit schemes need to be created to support solutions that lead to proven reductions in GHG emissions arising from fertilizer use.



## Responsible Plant Nutrition: $\uparrow$ NUE & $\downarrow$ N<sub>2</sub>O




**Figure 1.** Sources of nitrous oxide (N<sub>2</sub>O) emissions arising from nitrogen (N) inputs to agricultural soils. Emissions increase when fertilizers (F) or manures (M) are applied. Direct emissions occur when ammonium (NH<sub>4</sub><sup>+</sup>) is nitrified, and when nitrate (NO<sub>3</sub><sup>-</sup>) is denitrified. Indirect emissions arise from losses to drainage water and from ammonia (NH<sub>3</sub>) losses to air. Additional inputs include biological nitrogen fixation (B), atmospheric deposition (D), and irrigation water (W). The return of crop residues (CR) also contributes N to the soil pools from which N<sub>2</sub>O is emitted. Optimizing N use efficiency (NUE)—by matching inputs to crop removal (R) as closely as possible—is achieved through management of inputs (4R), crops and soils to optimize yields (Y). Improvements in NUE reduce fertilizer emissions per unit of crop produced. Nitrification inhibitors (NI), controlled release fertilizers (CRF), and urease inhibitors (UI) are effective 4R practices that decrease N<sub>2</sub>O emissions per unit of N applied.



# What is the issue?

Urgent action is required across all sectors of human activity to reduce GHG emissions and mitigate the impact of climate change, including the agrifood sector (1).

In 2021, emissions from the entire agri-food sector were about 16 billion tonnes of carbon dioxide equivalent (Gt CO<sub>2</sub>e), or about 30% of total anthropogenic emissions (53 Gt CO<sub>2</sub>e) (2). Agri-food emissions were largest in Asia (6.8 Gt CO<sub>2</sub>e) and the Americas (4.3 Gt CO<sub>2</sub>e), reflecting the large sizes of their populations and geographical areas. On-farm operations involved in producing food and other outputs (i.e. farm-gate) represented 48% of all agrifood systems emissions, while pre- (e.g. fertilizer production) and post-production (e.g. food packaging) processes contribute about 33% and land-use change 19% (Fig. 2). Due to increased agricultural productivity and efficiency, the per capita emission of GHG for food production has actually decreased over time. In 1990, the average person's food-related emissions amounted to 3.0 t CO<sub>2</sub>e/yr, but that number decreased to 2.4 t CO<sub>2</sub>e/yr by 2015 (3) and about 2 t CO<sub>2</sub>e/yr in 2021 per capita (2).



**Figure 2.** Annual greenhouse gas (GHG) emissions (Gt CO<sub>2</sub>e) of the agri-food sector by greenhouse gases and primary sources. Values shown are averages of 2017-2021. Values at the top of each bar are the sum of each component shown. The right-most column provides a breakdown of farm-gate N<sub>2</sub>O emissions. GHG emissions associated with production and transport of fertilizer (not shown) amount to ~0.5 of the 5.2 Gt CO<sub>2</sub>e shown for pre- & post-production. Source: FAOSTAT Emission totals.



## SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

The agrifood sector plays an important role in meeting global emission reduction targets (4), while it also has to ensure food security and soil health, preserve biodiversity, reduce environmental impacts, and improve farmers' incomes. Increasing crop yields and nutrient use efficiency on the existing agricultural land through responsible plant nutrition will be critical for achieving these goals (5). Providing nutrients in the form of mineral fertilizers to support crop growth results in GHG emissions associated with activities such as extraction, production, and processing. Emissions also arise from the transport and utilization at the farm level, i.e. from machinery used and from soils, the latter induced by application of N fertilizers (6).

Nitrogen is the essential plant nutrient with the largest GHG impact. Table 1 summarizes the global GHG emissions associated with the production and use of N fertilizers in 2018, adapted from a study which considered multiple sources of information, including the IPCC and FAO (7). Firstly, the energy-intensive process of synthetic N production is estimated to be responsible for 0.44 Gt or 440 Mt CO<sub>2</sub>e/yr. Secondly, N fertilizer input to soil increases microbial production of the powerful GHG nitrous oxide (N<sub>2</sub>O), termed direct N<sub>2</sub>O emissions; N losses in the form of ammonia volatilization and nitrate leaching further result in N<sub>2</sub>O emissions off-site, termed indirect N<sub>2</sub>O emissions (Fig. 1). Thus, the direct and indirect field emissions of N<sub>2</sub>O contributed 0.6 Gt CO<sub>2</sub>e of 52.7 Gt CO<sub>2</sub>e global emissions (Fig. 2) or 0.66 Gt CO<sub>2</sub>e if CO<sub>2</sub> release from urea is included (Table 1). Summing the production, transport, and in-field emissions, the synthetic N fertilizer supply chain contributed 1.13 Gt CO<sub>2</sub>e, or 2.1% of all global GHG emissions (Table 1). China, India, USA, and EU28 accounted for 62% of the total fertilizer N emissions. It should be noted, however, that these estimates have an average uncertainty of about ±15%, mostly due to uncertainties of direct and indirect N<sub>2</sub>O field emissions (±42 and ±24%, respectively) (7).

**Table 1.** Estimated global greenhouse gas emissions from production and use of inorganic N fertilizers in agriculture in 2018 (7). Note that the sum of indirect and direct N<sub>2</sub>O emissions equals 662 Mt CO<sub>2</sub>e (=0.662 Gt CO<sub>2</sub>e while values in Fig. 2 show 0.6 Gt CO<sub>2</sub>e due to the differences in the time period).

Fertilizer supply chain GHG emission components	Mt* CO <sub>2</sub> e	Share (%)
Synthetic N production	439	38.8
Transport	30	2.7
Direct field emissions (N <sub>2</sub> O, CO <sub>2</sub> )	466**	41.2
Indirect N <sub>2</sub> O emissions	196***	17.3
<b>Total emissions</b>	<b>1131</b>	<b>100</b>

\* million tons (1 Mt = 0.001 Gt)

\*\* 380 Mt CO<sub>2</sub>e as direct soil N<sub>2</sub>O emissions plus 86 Mt CO<sub>2</sub>e as CO<sub>2</sub> liberated from urea fertilizer

\*\*\* 66 Mt CO<sub>2</sub>e as indirect N<sub>2</sub>O derived from ammonia volatilization and redeposition and 130 Mt CO<sub>2</sub>e as indirect N<sub>2</sub>O derived from nitrate leaching



## SCIENTIFIC PANEL

### ON RESPONSIBLE PLANT NUTRITION

Emissions of N<sub>2</sub>O from manure management and application of other organic fertilizers add another 1.0 Gt CO<sub>2</sub>e annually (Fig. 2). Summing the above, direct and indirect N<sub>2</sub>O emissions associated with use of mineral N and organic fertilizers are in the order of about 2.13 Gt CO<sub>2</sub>e annually, or roughly 4% of the world's total emissions, representing a significant mitigation potential (Fig. 2). Producing and using other mineral fertilizers, such as phosphorus (P), or potassium (K), create additional GHG emissions that are not included here, but these have been estimated to be very small compared to N related emissions (e.g. 0.04 Gt CO<sub>2</sub>e for China in 2020 (8)).

While the use of N fertilizers contributes to GHG production, the efficient use of plant nutrients in agriculture can also play an important role in the mitigation of GHG emissions by:

1. Enabling crop production intensification on existing land, thus avoiding deforestation and land clearing which are the major causes of CO<sub>2</sub> emissions from land (see land-use change in Fig. 2)
2. Supporting the growth of biofuel crops as renewable energy sources
3. Supporting growth and removal of atmospheric CO<sub>2</sub> by plants that can lead to build up of soil organic matter (9)

Hence, opportunities to mitigate agri-food system GHG production include sustainably increasing crop yields, recycling more nutrients and increasing N use efficiency (NUE) on existing land, in addition to other actions such as prioritizing crop production for human consumption, changing diets, reducing food waste, and decarbonizing supply chains (10, 11). Importantly, increasing NUE and minimizing direct and indirect emissions of N<sub>2</sub>O present the most relevant opportunities for reducing emissions associated with fertilizers (6).

At issue is how to balance the need for using N fertilizers to support food production with the quest for reducing climate impacts of N fertilizer use, specifically by reducing direct and indirect N<sub>2</sub>O emissions. In this issue paper, we focus on the specific role of responsible plant nutrition to mitigate these GHG emissions. We discuss practices and technologies that mitigate GHG emissions related to N fertilizer use, and identify challenges to the implementation of solutions and the quantification of emission reductions. Finally, we suggest how various stakeholders can contribute to addressing these challenges.




SCIENTIFIC PANEL  
ON RESPONSIBLE PLANT NUTRITION

# What is happening?



## SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION



**Figure 3.** Changes in global cropland nitrogen (N) input, crop N yield, N surplus, nitrogen use efficiency (NUE) and farm gate nitrous oxide (N<sub>2</sub>O) emissions from 1990 to 2021. Source: FAOSTAT Emissions totals.

## Global Nitrogen Fertilizer Related Trends

The general global trajectory from the 1960s to 1990s has been an increase in N fertilizer use and crop production (N yield) accompanied by increasing N surplus and a decline or stagnation of cropland NUE (12). In the most recent decades, global crop production and N input (including fertilizer, manure, biological N fixation, seed, and atmospheric deposition) continued to rise. At the same time, global NUE increased to about 55% in 2021 (Fig. 3). The annual N surplus has remained fairly steady at about 80 million tons of N, although huge regional differences exist (3, 12). During this period, farm gate N<sub>2</sub>O emissions have risen to 2.1 Gt CO<sub>2</sub>e by 2021 (Fig. 3), mostly associated with mineral and organic fertilizer use (Fig. 2). The key challenge is to shift these trends towards more significant reductions of N surplus and N<sub>2</sub>O emissions – while continuing to increase productivity and NUE. The decrease in the ratio of N<sub>2</sub>O emission to N input apparent in Fig. 3 arises from an increase in the proportion of total N input from biological fixation, accompanied by a decrease in the proportion from manure.

## Characteristics of Nitrous Oxide Emissions

Nitrous oxide emission from soils primarily results from microbial nitrification and denitrification processes. These are regulated by numerous environmental conditions and specific abiotic processes (13). For example, after N fertilizer application and heavy rainfall events, a surplus of inorganic N and/or moisture in the soil enhances N<sub>2</sub>O emissions. Conditions that drive



## SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

N<sub>2</sub>O-producing microbial processes in soils are highly variable in space and time, resulting in emissions that are characterized as occurring in ‘hot spots’ and at ‘hot moments’, with a large percentage of emissions originating from brief, unpredictable events or from small areas in a field (14). The magnitude of these emission events is impacted by soil and climatic factors and their interactions with soil and crop management practices. The emission characteristics of soil N<sub>2</sub>O present challenges for measurement as well as identification and implementation of mitigation practices, which are generally regarded as being soil-, crop-, and climate-specific.





SCIENTIFIC PANEL  
ON RESPONSIBLE PLANT NUTRITION

# Interventions to reduce N<sub>2</sub>O emissions

A range of interventions have been shown to reduce soil N<sub>2</sub>O emissions, including the use of novel fertilizer products and improved agronomic practices (Fig. 1) (15).



## SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

These interventions in combination with soil carbon gains from improved productivity with balanced plant nutrition practices hold great promise as a means to reduce GHG emissions from agriculture (16). However trade-offs need to be considered as increasing soil organic carbon and hence soil fertility can increase N<sub>2</sub>O emissions, potentially offsetting climate change benefits from increased carbon sequestration (17). Nitrogen fertilizer additions need to be adjusted accordingly. Reducing N<sub>2</sub>O emissions is a permanent benefit in contrast to soil carbon gains which can be reversed with soil carbon release through subsequent agricultural practices such as increased tillage.

Effective interventions for soil N<sub>2</sub>O emission reduction are captured under the 4R Nutrient Stewardship framework which is focused on the 4 “rights” (source, rate, time and place) of nutrient applications. The New Paradigm for Responsible Plant Nutrition has broadened the scope of relevant practices and added to the 4R principles consideration of climate-smart fertilizers, i.e., using fertilizer sources with reduced carbon footprint, including emissions associated with both manufacture and use (18). Although users of the 4R framework often focus on mineral fertilizer use, its integration of organic fertilizer management is extremely important. For example, recycling organic N sources such as manure avoids GHG emissions associated with production of N fertilizer, while adjusting fertilizer N rate to account for N supplied by organic sources contributes to soil N<sub>2</sub>O emission reductions. The 4R practices discussed here need to be considered in a wider context of N management, including organic sources. A summary of the reductions in soil N<sub>2</sub>O emissions and changes in crop yield for various 4R and crop management practices is presented in Table 2. Owing to the high variability associated with N<sub>2</sub>O emissions, ‘summary effects’ of practices are typically derived from meta-analyses applied to multiple site-years of research across a wide range of conditions.

### • **Enhanced Efficiency Nitrogen Fertilizers**

Most meta-analyses confirm a strong effect of ‘enhanced-efficiency fertilizers’ (EEFs, including nitrification inhibitors, urease inhibitors and/or controlled-release fertilizers) on N<sub>2</sub>O emissions (Table 2). Consistent emission reductions are seen for these EEFs (36% across 608 peer-reviewed studies worldwide (19), but individual meta-analyses show a range of values for the different EEF types, with highest reductions generally observed from use of nitrification inhibitors (Table 2). In vegetable crops, controlled release fertilizers reduced N<sub>2</sub>O emissions on average by 24%, whereas nitrification inhibitors reduced them by 40%, with both technologies increasing yield by about 8% (20). Fertilizer sources can also effect N<sub>2</sub>O emissions in different ways. For example, in regions such as Ireland, which have mild, wet climates and high organic matter soils, significant reductions in N<sub>2</sub>O emissions were achieved by switching from nitrate-based fertilizers to urea with added urease or nitrification inhibitors (21). On the other hand, in a field experiment with drip-fertigated crops in Spain, lower emissions were measured from calcium nitrate as compared to urea (22).

In summary: (i) reductions in N<sub>2</sub>O emissions due to EEFs are generally larger than their positive effects on crop yield or NUE (23)); (ii) nitrification inhibitors are most effective in reducing N<sub>2</sub>O emissions, but may also increase ammonia volatilization losses; and (iii) in many situations combining urease and nitrification inhibitors can be recommended to avoid pollution swapping. For example, in the case of using a urease inhibitor alone, direct N<sub>2</sub>O emissions may increase even though indirect N<sub>2</sub>O emissions may be reduced (24, 25). In addition to impacts on GHG emissions, EEFs generally also reduce nitrate leaching losses from the soil by about 20-40% (26). This decrease in N loss provides additional



**SCIENTIFIC PANEL**

ON RESPONSIBLE PLANT NUTRITION

environmental and economic benefits, through reduced indirect N<sub>2</sub>O emissions, reduced contamination of groundwater, and reduced fertilizer costs.

- Fertilizer Timing and Placement**

The effects of timing and placement of N fertilizers are usually smaller and less consistent than those of EEF (Table 2). Young et al. (19), in a synthesis of 113 meta-analysis studies, reported that optimal fertilizer placement decreased direct N<sub>2</sub>O emissions by 15% and NH<sub>3</sub> emissions by 43% (which would reduce indirect N<sub>2</sub>O emissions), and increased yield by 5%. Optimal timing had no significant effect on direct N<sub>2</sub>O emissions, while on average it reduced NH<sub>3</sub> emissions by 32% (Table 2).

*\* n represents number of meta-analyses for source (19) and number of site-years for the rest*

*\*\*EEF: enhanced-efficiency fertilizers that included nitrification/urease inhibitors and slow-release fertilizers*

**Table 2.** Relative changes (%) in N<sub>2</sub>O and NH<sub>3</sub> emissions and crop yield due to 4R and crop management practices. Sources: Meta-analyses summarizing the global literature.

Practice	N <sub>2</sub> O emission %	NH <sub>3</sub> emission %	Crop yield %	n* (for N <sub>2</sub> O)	Source
Crop rotation	-4		-2	1	(19)
Cover cropping	-5		12	3	(19)
Fertilizer rate	-9	-31	-4	7	(19)
Fertilizer timing	0	-32	4	3	(19)
Fertilizer placement	-15	-43	5	1	(19)
Fertilizer source:					
Nitrification inhibitor	-50	20	2	422	(27)
Urease + nitrification inhibitor	-30	-38	3	118	(27)
Urease inhibitor	-10	-50	6	116	(27)
Controlled-release fertilizer	-19 to -33	-65	-3	276	(28–30)
Maize EEF**	-25		4	429	(30, 31)
Wheat EEF**	-29		5	279	(30, 31)
Rice EEF**	-25		8	312	(30, 31)



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

- **Biological Nitrification Inhibitors**

While the studies on inhibitors summarized in Table 2 refer only to manufactured inhibitors added to inorganic fertilizers, there also exists the potential to modify biological nitrification based on compounds exuded by plants that may suppress nitrification and N<sub>2</sub>O emission. A recent survey of the literature found that “Primary metabolites, such as sugars, amino acids, and organic acids, strongly stimulated soil N<sub>2</sub>O emissions, by an average of 79%, while secondary metabolites, such as phenolics, terpenoids, and flavonoids, often characterized as both biological nitrification inhibitors (BNIs) and biological denitrification inhibitors (BDIs), reduced soil N<sub>2</sub>O emissions by an average of 41%.” (32). Biological nitrification inhibition explains, for example, the low N<sub>2</sub>O emission rates observed from *Brachiaria*, a grass species often used as pastures and cover crops in Brazil. Some potential may exist to strengthen BNI traits in high-yielding cereal crops through plant breeding (33, 34). However, feasibility and efficacy of BNI compared to synthetic nitrification inhibitors has not been fully explored.

- **Nitrogen Rate**

The rate of N fertilizer applied has been widely used in GHG accounting to estimate N<sub>2</sub>O emissions. The IPCC established an average fertilizer-induced direct emission factor (EF) of 1% (uncertainty range: 0.1-1.8%) for global estimates; that is, for every 100 kg N applied, 1 kg of N<sub>2</sub>O-N is emitted directly from soils (35). The IPCC provides some further differentiation, e.g. a 1.6% EF for fertilizer used in ‘wet’ climates, or an EF of only 0.3% in continuously flooded rice fields. Although the IPCC value has proven robust for N<sub>2</sub>O emission estimates for regions that lack locally derived EFs, research has shown that EFs can vary widely and, in many cases, are also significantly lower than 1%. In Australia, for example, the average EF of mineral N fertilizer was 0.70%, with a range of 0.17-1.77% across regions and cropping systems (36). In Germany, the district-wise EFs ranged from 0.38% to 0.92%, with a national average of 0.62% (37). In a global spatial analysis, average estimated EFs were 1.02% for maize, 0.58% for wheat, and 0.52% for rice, but varied widely, including hotspots with higher values (38). Similarly, Wang et al. (39) generated global maps of EFs with an average of 0.88% for maize and 0.65% for wheat. Yao et al. (31) found average global EFs of 0.69% for maize, 0.60% for wheat, and 0.36% for rice.



## SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

On the other hand, the global agri-food system also includes production systems in which N application rates, N<sub>2</sub>O emissions, and EFs are particularly large. For example, although tea plantations account for only 0.3% of total cropland area, they have been estimated to account for 1.5%–12.7% of total direct cropland N<sub>2</sub>O emissions, with average losses of 17 kg N ha<sup>-1</sup> in the form of N<sub>2</sub>O emissions and a global mean EF of 2.3% (40). These high EFs are driven by N application rates that far exceed the tea crop N demand, inducing an exponential increase in soil N<sub>2</sub>O emissions with rates exceeding 250 kg N ha<sup>-1</sup> yr<sup>-1</sup>. Such global hotspots must be addressed with particular urgency.

The general assumption in using EFs is that the N<sub>2</sub>O emission rate is a linear function of N rate, an assumption that is likely incorrect. Shcherbak et al. (41) reviewed 233 site-years of studies with at least three N levels and found that N<sub>2</sub>O emissions responded non-linearly to N rate, demonstrating that reducing N that is in surplus of crop needs would be most effective in reducing emissions. The principles of plant nutrition suggest that at the optimal N rate for a crop grown in a specific environment, NUE is high, N surplus is low, and N losses and N<sub>2</sub>O emissions are also likely to be lower than at rates exceeding the optimum. This principle has been implemented, for example, in the N balance model of the Environmental Defence Fund (42, 43). The key implication is that regions where N is applied in excess of optimal fertilizer rates have the largest N<sub>2</sub>O mitigation potential arising from rate reduction.





SCIENTIFIC PANEL  
ON RESPONSIBLE PLANT NUTRITION

# Interventions to increase yields and N use efficiency

In regions with low N use or nutrient deficiency there is little potential to mitigate current N<sub>2</sub>O emissions, but the need to increase yields on existing cropland to improve food security and avoid the CO<sub>2</sub> emissions that arise from deforestation and land clearing must be recognized.



## SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

These regions are particularly common in sub-Saharan Africa. Many of these regions feature soils that have been degraded and require both mineral fertilizers and organic inputs, along with sound agronomic practices, superior seed varieties, soil amendments, and other measures, as prescribed by the principles of Integrated Soil Fertility Management, or ISFM (44). There are also opportunities to sequester carbon in these soils (45, 46), but such carbon first needs to be produced through increased crop yields.

The assessment of the full net impacts on GHG emissions, land-use change, and soil carbon sequestration arising from the wide array of site-specific ISFM practices is beyond the scope of this paper. It should nevertheless be recognized that the simultaneous improvements achieved in yields and N use efficiency in North America or other regions depended in large part on massive investments in crop genetic improvement as well as soil, crop and nutrient management (47, 48). Similar investments will likely be required for other parts of the world where crop yields are far below potential, particularly in Sub-Saharan Africa (49).

## Summary

A wide range of agronomic practices and 4R N management solutions have been shown to increase NUE and/or directly reduce N<sub>2</sub>O emissions across regions and crops. These effective actions have potential to mitigate a large fraction of the 0.66 Gt CO<sub>2</sub>e annual emissions of GHG from fertilizer use within the next few decades, particularly when applied in combination (6, 39). These actions would make an important contribution towards achieving net-zero emissions by 2050 by reducing global agri-food system emissions (16 Gt CO<sub>2</sub>e) (4). Due to their large area and share of global N fertilizer consumption, grain production systems play a particularly important role in implementing GHG mitigation actions. At issue is how to achieve large-scale implementation of such measures.



SCIENTIFIC PANEL  
ON RESPONSIBLE PLANT NUTRITION

# What can be done?



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

## Recommended Implementation Actions

Identifying the most promising practices to reduce N<sub>2</sub>O emissions is a necessary first step to achieve reduction targets but, ultimately, implementation is dependent on buy-in, trust and actions from farmers. Depending on the region, practices that are ‘win-win’ may already be on their way to being adopted, or potential exists for adoption with appropriate knowledge and technology transfer programs. For other GHG reduction practices and technologies, incentives for farmers to adopt them could include reduced input costs, improved productivity, subsidized EEF products, payments, and sustainability-linked finance. Improved agronomic practices that boost crop productivity with the minimum amount of N fertilizer needed can incentivize farmers by helping them reduce input costs, while increasing income.

We group the potential solutions into three action targets, which are outlined in Table 4. Regions in which excessive N rates result in substantial N surplus should focus first on target 1 practices. Examples include many parts of China and India, but also some areas in Europe and North America with high cropping and livestock intensity (50). In regions with substantial N surplus, the first priority would be to deploy the Right Rate principle, providing a win-win situation since GHG emissions would be reduced while profits would be increased. The achievable reductions could be significant. However, it is important that a non-linear EF model be used to capture decreases in N<sub>2</sub>O emissions from reductions in N surplus as the linear EF model would underestimate potential emission reductions (41).





**Table 4.** Overview of broad actions targeting N<sub>2</sub>O emission reductions and their potential benefits

Action Target	Practices	Benefits
1. Reduce excessive surplus nitrogen.	Right Rate, achieved through improved crop N demand prediction and improved manure management, and agronomic management to improve crop yield and N uptake.	Direct and indirect N <sub>2</sub> O emissions are reduced; yield is not affected; NUE and profits are increased.
2. Reduce ammonia losses.	Right Time, Right Placement, and use of urease inhibitors or controlled-release fertilizers (Right Source) with adjustments in rate.	Indirect N <sub>2</sub> O emissions are reduced; yield and NUE are increased; profits are potentially increased if rate is adjusted.
3. Reduce direct N <sub>2</sub> O losses.	Use of nitrification inhibitors and controlled-release fertilizers (Right Source) with the Right Rate, Right Time and Right placement.	GHG emissions are reduced; yield and NUE are not affected or may increase slightly.

In regions where fertilizer N rates are already close to optimal (Right Rate), the focus should be on reducing N loss as ammonia through better timing, placement, and use of urease inhibitors or controlled-release fertilizers (Target 2; Table 4). Ammonia emissions can comprise a substantial fraction of the fertilizer N applied as urea. Reducing emissions would decrease indirect N<sub>2</sub>O emissions and result in significantly increased yields, NUE, and profits. However, care must be taken to adjust the N rate to account for the ammonia reductions; otherwise, direct N<sub>2</sub>O emissions can be increased due to higher N retention in the soil.

The third target would apply to situations where yields and N use efficiency are already close to optimal. Actions would prioritize direct N<sub>2</sub>O emission reduction through practices such as using controlled-release fertilizers, nitrification inhibitors, or combinations of nitrification and urease inhibitors. Because N<sub>2</sub>O emissions are small compared to N fertilizer input, nitrification inhibitors may not result in substantial yield increases and may not increase profits (Table 2). For that reason, support mechanisms may often be required to incentivize target 3 action. All three target practices could also reduce nitrate leaching, and thus have additional environmental benefits. It is likely that N loss in dinitrogen (N<sub>2</sub>) form through denitrification would also be mitigated with target 3 actions, providing economic benefits, but there is little evidence for this given the difficulty in measuring N<sub>2</sub> gas emissions.

A combination of 4R practices is likely needed to achieve the largest GHG reductions. The proposed action categories in Table 4 are meant to be cumulative. For example, there may still be opportunity to adjust N fertilizer rates for areas that fall under targets 2 and 3 where rates are not excessive and close to the optimum due to uncertainty in crop N demand predictions. In these cases, farmers may still tend to use slightly more N fertilizer than needed to reach the economic optimum and additional incentives related to risk management may be required.



## Policies to Drive Implementation

The measures listed in Table 4 are potential means to improve NUE and reduce GHG emissions as well as other forms of N pollution, while meeting food security goals. Hence, redirecting public policy and subsidies towards their wide adoption should result in carefully designed regulations or incentives that are cost-effective, target hotspots and consider the trade-offs among yield, profit, and environmental footprint (51). Nitrogen use efficiency improvement targets can play a central role in achieving the right balance among the different goals to pursue. In China, for example, stricter environmental protection policies and improved N management led to a nationwide reduction in N fertilizer applied per area, increased NUE and slowed down growth in cropland N<sub>2</sub>O emissions after 2003 (52, 53).

Overall, however, evidence for successful policies that incentivize NUE increases, reductions in N pollution and N<sub>2</sub>O emissions while also supporting food production remains quite scarce (54). For a long time, integrating N<sub>2</sub>O mitigation into climate policy has received too little attention (55). It should also be noted that further reductions in N loss in the long term will require profound food system changes, particularly shifts in diets in regions with high levels of food security (56).

Incentives to adopt N<sub>2</sub>O emission reduction measures can also come from carbon markets that would reward farmers financially for implementing practices to reduce emissions. Carbon credits in this case can be associated with any of the actions listed in Table 4, i.e. reducing the application rate of inorganic and organic fertilizers, increasing NUE, and/or directly reducing N<sub>2</sub>O emissions through inhibitors or controlled-release fertilizers. Evidence-based methodologies (standards) must still be established for such purposes,



## SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

including monitoring and verification. There is also growing interest in addressing Scope 3 GHG emissions by companies providing services or procuring products from the agricultural sector. Scope 3 emissions “are the result of activities from assets not owned or controlled by the reporting organization, but that the organization indirectly affects in its value chain” (57). On-farm emissions such as soil N<sub>2</sub>O emissions are significant Scope 3 emissions for companies selling N fertilizer or for companies purchasing agricultural products to be used within their supply chain. For example, a recent partnership between Nutrien Ag Solutions and Maple Leaf Foods for 4R Stewardship adoption by farmers in the Canadian Prairies is a value chain intervention addressing Scope 3 emissions by both companies. It aims to produce food products with lower carbon intensity by growing crops (e.g., canola, barley, wheat, peas) with optimized N fertilizer management (58).

Sustainability finance may also incentivize low-emission practices. Governments, for example, may re-direct some of their financial support to farmers adopting NUE-increasing and emission-reducing practices. In other cases, banks may provide improved terms of finance, such as lower interest rates, as farm environmental performance improves. Such policy and market mechanisms may provide the needed incentives to tackle in particular target 3 type emission reduction practices (Table 4) which benefit the public more than the farmer.

## Metrics to Monitor Progress

Regardless of the mechanism used to incentivize action, metrics to monitor progress towards reduction or efficiency goals are needed. Sustainability metrics are an essential element of tracking the impact of improved N fertilizer management and to determine efforts by farmers, companies, or government in reducing their carbon footprints and meeting emission reduction targets. The availability of metrics also limits which practices are eligible for incentives. Existing GHG methodologies are mostly based on IPCC guidelines that only consider the N rate, but not other 4R practices such as placement or use of inhibitors. Hence, carbon reduction protocols often focus on reducing the N rate rather than increasing NUE, which may create undesirable yield losses too.

An exception is the Nitrous Oxide Emission Reduction Protocol (NERP) implemented in Alberta, Canada, which included 4R practices beyond N rate (59). The approach was adopted by several supply-chain initiatives which aim to quantify and monitor GHG emissions from organizations along the agricultural value chain. However, several gaps remain, including evaluating regionally specific management practices, assessing the 4Rs used as suites of practices rather than individually, evaluating variable-rate and source fertilizer technology, and quantifying the impact of single vs. dual use of urease and nitrification inhibitors. Furthermore, the interactions with other conservation practices, such as diversifying crop rotations or cover crops are not represented yet.



## SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

Additionally, a pragmatic approach could be to estimate N<sub>2</sub>O emissions as a function of the partial crop N balance. As suggested by several studies, this would include the difference between N inputs (fertilizer, manure, other sources) and N removed by crop harvest (15, 43, 60, 61). Emissions of N<sub>2</sub>O have been found to relate as well or better to N balance than to the rate of N applied (15). Generalized relationships have been developed for multiple crops in North America (43) and globally (15). These show that, in cereal cropping systems, N<sub>2</sub>O emissions are likely to be higher once the N surplus exceeds about 50 kg N/ha. Using surplus reduction as an indicator of emissions is relatively easy to implement in farming through farm management software. Similarly, the information collected would allow monitoring NUE and N surplus as generally important indicators for agronomic and environmental farm performance, including guidance on targets to achieve (62). A focus on surplus, however, does not capture all possible reductions in emissions associated with fertilizer, particularly occurring as short-lived peaks or in hotspots. It also neglects independent effects of inhibitors and controlled-release products on direct and indirect N<sub>2</sub>O emissions, as noted above.

A final consideration is about the way to express emissions (and their reductions). Some metrics typically express emissions per unit of land area (kg N<sub>2</sub>O-N per ha), as a GHG intensity (per unit of crop production, e.g. N<sub>2</sub>O-N per kg grain) or per unit of N input provided (i.e. as an emission factor, kg N<sub>2</sub>O-N per kg N applied). It is likely that all three metrics are useful depending on the aim of the emission reduction. GHG intensity and EFs are efficiency-based metrics which account for output or input, while a land-based metric does not consider the input received or output provided by the land area. For absolute emission reductions, land-based emissions are the most relevant metric. In carbon credit

schemes for food supply chains, however, companies wanting to reduce the carbon footprint may prefer using per-unit-of-food (intensity-based) measures. Emissions per unit of product are also important for international trade, better suiting the application of carbon border adjustments.

Advances in N<sub>2</sub>O emission knowledge and practices to reduce emissions indicate that mitigation following the principles of Responsible Plant Nutrition is possible and feasible. To advance implementation at the farm level, incentive mechanisms have to recognize motivations and barriers to adoption and prioritize actions. Making this happen will require commitments and aligned actions by many stakeholders.



# Who needs to do what?

## Governments:

- Adopt policies that incentivize and reward adoption of practices and technologies that have proven impact on increasing N use efficiency and reducing GHG emissions, N<sub>2</sub>O in particular, such as nitrification and urease inhibitors and controlled-release N fertilizers.
- Work with industry on last-mile delivery extension programs and emission-reporting regulations compatible with market-based approaches.
- Improve methodologies and ensure that reduction efforts are captured in national GHG inventory reports.

## Industry:

- Implement targeted Scope 3 emissions reduction programs; incentivize and support the documentation of emission reductions.
- Adopt standards for reducing N<sub>2</sub>O emissions and join voluntary carbon market schemes.
- Support and train accredited crop advisers to facilitate extension, monitoring, reporting and verification of emission-reducing practices.
- Drive innovation, as well as provide resources to develop new technologies for GHG mitigation, including smart, controlled-release fertilizers, and real-time precision N management technologies.
- On a regional basis, provide leveraged support for public research on key issues related to the 4Rs.

## Carbon market participants and investors:

- Develop and refine evidence-based, transparent standards for claiming carbon credits through adoption of N<sub>2</sub>O emission-reducing practices.
- Create and implement voluntary carbon market schemes that financially benefit farmers and stimulate wider adoption of practices and technologies that reduce N<sub>2</sub>O emissions.



# Who needs to do what?

## Farmers and other practitioners:

- Optimize nutrient management with available 4R practices based on crop needs and soil fertility conditions, and using appropriate decision support tools.
- Evaluate practices at the farm level; keep records and provide data on farm activities that can be used in protocols and sustainability metrics; provide technical assistance (peer-to-peer).

## Researchers:

- Design and evaluate novel practices that increase NUE and reduce N<sub>2</sub>O emissions.
- Develop and improve databases, emission factors, methodologies, and models for use in emission reduction protocols and standards.
- Evaluate cost/benefits of practices and work with farmers/practitioners to identify practices that are most feasible and practical at the farm level.



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

# What would success look like?

If the actions outlined in Table 4 are adopted widely enough, the following results could be achieved in the coming few decades:

## 1

Approximately 70% of the GHG emissions associated with fertilizer use could be mitigated by 2050. The bulk of that must be achieved in regions that account for most of the current N fertilizer use: particularly Asia, North America, and Europe.

## 2

The IPCC methodology for GHG emission inventories estimates N<sub>2</sub>O emissions based on N balance rather than N rates, recognizing that N rate increases in regions of low N use increase emissions less than in regions with high rates of N use.

## 3

The wide adoption of practices that reduce N<sub>2</sub>O emissions also supports further increases in crop yields and N use efficiency and reduces other losses of reactive N to the environment, thus contributing to global ambitions of halving N waste, improving water quality and protecting biodiversity.

## 4

Robust methods and standards have been widely adopted to account for N<sub>2</sub>O emissions in farming, and to support carbon credit schemes that financially incentivize N<sub>2</sub>O-reducing technologies and practices. Farmers gain from these new payment schemes and can offset costs associated with adopting new technologies and practices. Mainstreaming incentivizes industry and innovators to invest in developing even better technologies and solutions.



# What would success look like?

**5**

Participation by farmers in agri-food industry schemes to reduce the GHG footprint of food has led to more trust and transparency, including willingness to share data on nutrient management practices for purposes of monitoring, evaluation, and reporting of sustainability outcomes. Digital technologies are increasingly used for these purposes, including artificial intelligence.

**6**

Along with decarbonized fertilizer production, new product and food labels have emerged that report their full-chain carbon footprints inclusive of the portion associated with fertilizers.

**7**

Researchers and startups have developed novel fertilizers with minimal GHG emissions and longer-lasting modes of action for specific conditions.



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

## References

1. H. Tian et al., A comprehensive quantification of global nitrous oxide sources and sinks. *Nature*. 586, 248–256 (2020).
2. FAO, Agrifood systems and land-related emissions. Global, regional and country trends, 2001–2021. FAOSTAT Analytical Briefs Series No. 73 (2023).
3. M. Crippa et al., Food systems are responsible for a third of global anthropogenic GHG emissions. *Nature Food*. 2, 198–209 (2021).
4. C. Costa et al., Roadmap for achieving net-zero emissions in global food systems by 2050. *Sci Rep*. 12, 15064 (2022).
5. A. Dobermann et al., Responsible plant nutrition: A new paradigm to support food system transformation. *Global Food Security*. 33, 100636 (2022).
6. IFA & Systemiq, Reducing emissions from fertilizer use (International Fertilizer Association and SystemIQ, Paris, France, 2022).
7. S. Menegat, A. Ledo, R. Tirado, Greenhouse gas emissions from global production and use of nitrogen synthetic fertilisers in agriculture. *Sci Rep*. 12, 14490 (2022).
8. H. Gong et al., Synergies in sustainable phosphorus use and greenhouse gas emissions mitigation in China: Perspectives from the entire supply chain from fertilizer production to agricultural use. *Science of the Total Environment*. 838, 155997 (2022).
9. J. W. van Groenigen et al., Sequestering soil organic carbon: a nitrogen dilemma. *Environmental Sci. & Technol*. 51, 4738–4739 (2017).
10. J. D. Gil et al., Reconciling global sustainability targets and local action for food production and climate change mitigation. *Global environmental change : human and policy dimensions*. 59, 101983 (2019).
11. A. Muscat et al., Principles, drivers and opportunities of a circular bioeconomy. *Nature Food*. 2, 561–566 (2021).
12. X. Zhang et al., Managing nitrogen for sustainable development. *Nature*. 528, 51–59 (2015).
13. K. Butterbach-Bahl et al., Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philosophical Trans. Royal Society of London. Series B, Biological sciences*. 368, 20130122 (2013).
14. C. Wagner-Riddle et al., Mitigation of nitrous oxide emissions in the context of nitrogen loss reduction from agroecosystems: managing hot spots and hot moments. *Current Opinion in Environmental Sustainability*. 47, 46–53 (2020).
15. T. M. Maaz et al., Meta-analysis of yield and nitrous oxide outcomes for nitrogen management in agriculture. *Global Change Biol*. 27, 2343–2360 (2021).
16. N. C. Lawrence, C. G. Tenesaca, A. VanLoocke, S. J. Hall, Nitrous oxide emissions from agricultural soils challenge climate sustainability in the US Corn Belt. *Proc. Natl. Acad. Sci. U. S. A.* 118 (2021).
17. B. Guenet et al., Can N<sub>2</sub>O emissions offset the benefits from soil organic carbon storage? *Global Change Biol*. 27, 237–256 (2021).
18. Scientific Panel on Responsible Plant Nutrition, Furthering 4R nutrient stewardship. Issue Brief 03 (SPRPN, Paris, France, 2022).
19. M. D. Young, G. H. Ros, W. de Vries, Impacts of agronomic measures on crop, soil, and environmental indicators: A review and synthesis of meta-analysis. *Agriculture, Ecosystems & Environment*. 319, 107551 (2021).
20. Z. Pan et al., Global impact of enhanced-efficiency fertilizers on vegetable productivity and reactive nitrogen losses. *Science of the Total Environment*. 926, 172016 (2024).



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

## References

21. M. A. Harty et al., Reducing nitrous oxide emissions by changing N fertiliser use from calcium ammonium nitrate (CAN) to urea based formulations. *Science of the Total Environment*. 563-564, 576–586 (2016).
22. D. Abalos et al., Management of irrigation frequency and nitrogen fertilization to mitigate GHG and NO emissions from drip-fertigated crops. *Science of the Total Environ.* 490, 880–888 (2014).
23. D. Abalos et al., Meta-analysis of the effect of urease and nitrification inhibitors on crop productivity and nitrogen use efficiency. *Agriculture, Ecosystems & Environment*. 189, 136–144 (2014).
24. C. F. Drury et al., Combining urease and nitrification inhibitors with incorporation reduces ammonia and nitrous oxide emissions and increases corn yields. *J. Environ. Qual.* 46, 939–949 (2017).
25. A. L. Woodley et al., Ammonia volatilization, nitrous oxide emissions, and corn yields as influenced by nitrogen placement and enhanced efficiency fertilizers. *Soil Sci. Soc. Am. J.* 84, 1327–1341 (2020).
26. FFAR, The Global Fertilizer Challenge: Future The Global Fertilizer Challenge: directions for efficient fertilizer research (2023).
27. M. Chen et al., Evidence map of the benefits of enhanced-efficiency fertilisers for the environment, nutrient use efficiency, soil fertility, and crop production. *Environ. Res. Lett.* 18, 43005 (2023).
28. R. Thapa et al., Effect of enhanced efficiency fertilizers on nitrous oxide emissions and crop yields: a meta-analysis. *Soil Sci. Soc. Am. J.* 80, 1121–1134 (2016).
29. B. Pan et al., Ammonia volatilization from synthetic fertilizers and its mitigation strategies: A global synthesis. *Agriculture, Ecosystems & Environment*. 232, 283–289 (2016).
30. D. Grados et al., Synthesizing the evidence of nitrous oxide mitigation practices in agroecosystems. *Environ. Res. Lett.* 17, 114024 (2022).
31. Z. Yao et al., A global meta-analysis of yield-scaled N<sub>2</sub>O emissions and its mitigation efforts for maize, wheat, and rice. *Global Change Biol.* 30, e17177 (2024).
32. Y. Lu et al., Biological mitigation of soil nitrous oxide emissions by plant metabolites. *Global Change Biol.* 30, e17333 (2024).
33. G. V. Subbarao, T. D. Searchinger, A "more ammonium solution" to mitigate nitrogen pollution and boost crop yields. *Proc. Natl. Acad. Sci. U. S. A.* 118, e2107576118 (2021).
34. G. V. Subbarao et al., Enlisting wild grass genes to combat nitrification in wheat farming: A nature-based solution. *Proc. Natl. Acad. Sci. U. S. A.* 118, e2106595118 (2021).
35. IPCC, 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (2019).
36. P. Grace et al., Revised emission factors for estimating direct nitrous oxide emissions from nitrogen inputs in Australia's agricultural production systems: a meta-analysis. *Soil Res.* 62 (2024).
37. G. P. Mathivanan et al., New N<sub>2</sub>O emission factors for crop residues and fertiliser inputs to agricultural soils in Germany. *Agriculture, Ecosystems & Environment*. 322, 107640 (2021).
38. X. Cui et al., Global mapping of crop-specific emission factors highlights hotspots of nitrous oxide mitigation. *Nat. Food.* 2, 886–893 (2021).
39. C. Wang et al., Reducing soil nitrogen losses from fertilizer use in global maize and wheat production. *Nat. Geosci.*, 1–8 (2024).
40. Y. Wang et al., Tea-planted soils as global hotspots for N<sub>2</sub>O emissions from croplands. *Environ. Res. Lett.* 15, 104018 (2020).
41. I. Shcherbak, N. Millar, G. P. Robertson, Global metaanalysis of the nonlinear response of soil nitrous oxide (N<sub>2</sub>O) emissions to fertilizer nitrogen. *Proc. Natl. Acad. Sci.* 111, 9199–9204 (2014).
42. EDF, How to use EDF's nitrogen balance model to make nitrous oxide and nitrate reduction claims (2022).



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

## References

43. A. J. Eagle et al., Quantifying on-farm nitrous oxide emission reductions in food supply chains. *Earth's Future*. 8 (2020).
44. Scientific Panel on Responsible Plant Nutrition, Fertilizer and soil health for enhanced productivity and sustainability in sub-Saharan Africa. Issue Brief 06 (SPRPN, Paris, France, 2024).
45. M. Lessmann et al., Global variation in soil carbon sequestration potential through improved cropland management. *Global Change Biol.* 28, 1162–1177 (2022).
46. R. Lal, Carbon farming by recarbonization of agroecosystems. *Pedosphere*. 33, 676–679 (2023).
47. T. Fischer et al., Sixty years of irrigated wheat yield increase in the Yaqui Valley of Mexico: Past drivers, prospects and sustainability. *Field Crops Res.* 283, 108528 (2022).
48. S. M. Mueller, C. D. Messina, T. J. Vyn, Simultaneous gains in grain yield and nitrogen efficiency over 70 years of maize genetic improvement. *Sci. Rep.* 9, 9095 (2019).
49. B. Vanlauwe et al., Fertilizer and soil health in Africa: The role of fertilizer in building soil health to sustain farming and address climate change (2023).
50. L. F. Schulte-Uebbing et al., From planetary to regional boundaries for agricultural nitrogen pollution. *Nature*. 610, 507–512 (2022).
51. G. Mandrini et al., Exploring trade-offs between profit, yield, and the environmental footprint of potential nitrogen fertilizer regulations in the US Midwest. *Front. Plant Sci.* 13, 852116 (2022).
52. Xuejun LIU et al., A new approach to holistic nitrogen management in China. *Front. Agr. Sci. Eng.* 0, 0 (2022).
53. Z. Shang et al., Weakened growth of cropland-N<sub>2</sub>O emissions in China associated with nationwide policy interventions. *Global Change Biol.* 25, 3706–3719 (2019).
54. D. R. Kanter et al., Gaps and opportunities in nitrogen pollution policies around the world. *Nat. Sustain.* 3, 956–963 (2020).
55. D. R. Kanter, S. M. Ogle, W. Winiwarter, Building on Paris: integrating nitrous oxide mitigation into future climate policy. *Current Opinion in Environmental Sustainability*. 47, 7–12 (2020).
56. A. Leip et al., Halving nitrogen waste in the European Union food systems requires both dietary shifts and farm level actions. *Global Food Security*. 35, 100648 (2022).
57. USEPA, Scope 3 inventory guidance (2024).
58. SustainCert, Nutrien/Maple Leaf Foods - Reduced N<sub>2</sub>O emissions from 4R Nitrogen Management (2024).
59. Government of Alberta, Quantification protocol for agricultural nitrous oxide emission reductions. Version 2.1 (2023).
60. Scientific Panel on Responsible Plant Nutrition, Defining nutrient use efficiency in responsible plant nutrition. Issue Brief 04 (SPRPN, Paris, France, 2023).
61. E. L. McLellan et al., The nitrogen balancing act: tracking the environmental performance of food production. *BioScience*. 68, 194–203 (2018).
62. EU Nitrogen Expert Panel, Nitrogen Use Efficiency (NUE). Guidance document for assessing NUE at farm level (Wageningen University, Alterra, Wageningen, NL, 2016).



SCIENTIFIC PANEL

ON RESPONSIBLE PLANT NUTRITION

## Authors, citation and contact

**Authors:** Members of the Scientific Panel on Responsible Plant Nutrition (lead authors underlined).

Patrick H Brown (University of California-Davis, USA), Tom Bruulsema (Plant Nutrition Canada), Achim Dobermann (International Fertilizer Association, France), Mariangela Hungria (Embrapa, Brazil), Jian Feng Ma (Okayama University, Japan), Michael McLaughlin (University of Adelaide, Australia), Bernard Vanlauwe (International Institute of Tropical Agriculture, Kenya), Marta Vasconcelos (Universidade Católica Portuguesa, Portugal), Claudia Wagner-Riddle (University of Guelph, Canada), Fangjie Zhao (Nanjing Agricultural University, China)

**Citation:** Scientific Panel on Responsible Plant Nutrition. 2024. Mitigating greenhouse gas emissions through responsible plant nutrition. Issue Brief 07. Available at <https://sprpn.org>.

**Further information:** Scientific Panel on Responsible Plant Nutrition, [info@sprpn.org](mailto:info@sprpn.org)